

Cincinnati, Ohio - 2006 (N=300)

Figure A. Age of GISP participants, in years, 2006




Figure B. Race/ethnicity of GISP participants, 2006




Figure C. Percentage of GISP participants identifying as men who have sex with men, 1988-2006






Figure D. Reason for visit among GISP participants, 2006



Cincinnati, Ohio - 2006 (N=300)

Figure E. Previous episode of gonorrhea among GISP participants, 1991-2006



*Data first collected in 1991. **Data first collected in 1992.

Note: Data points not shown when >30% data missing.

Figure F. Drugs used to treat gonorrhea among GISP participants, 2006




Figure G. Drugs used to treat *Chlamydia trachomatis* infection among GISP participants, 2006





Figure H. Resistance to penicillin, tetracycline, and ciprofloxacin among GISP isolates, 2006



Cincinnati, Ohio - 2006 (N=300)

Figure I. Decreased susceptibility to ceftriaxone among GISP isolates, 1990-2006




Figure J. Decreased susceptibility to cefixime among GISP isolates, 1992-2006






Figure K. Intermediate resistance and resistance to ciprofloxacin among GISP isolates, 1990-2006



Note: Susceptibility to ciprofloxacin first measured in 1990.

Figure L. Decreased susceptibility to azithromycin among GISP isolates, 1992-2006



Note: Susceptibility to azithromycin first measured in 1992.

Note: Decreased susceptibility to azithromycin is defined here as $\geq 1.0 \mu\text{g/ml}$.
No NCCLS criteria currently exist.